

Papers in other journals

Compiled by Becky Aston, Rosie Butler, Gill Brook, Jo Dafforn, Roberta Eales, Emma Hammerton and Claire-Anne Head

Obstetrics

- Altman D., Ekström Å., Forsgren C., Nordenstam J. & Zetterström J. (2007) Symptoms of anal and urinary incontinence following cesarean section or spontaneous vaginal delivery. *American Journal of Obstetrics and Gynecology* **197** (5), 512.e1–512.e7.
- Bastiaenen C. H., de Bie R. & Essed G. (2007) Pregnancy-related pelvic girdle pain. *Acta Obstetricia et Gynecologica Scandanavica* **86** (10), 1277–1278.
- Bellamy L., Casas J.-P., Hingorani A. D. & Williams D. J. (2007) Pre-eclampsia and risk of cardiovascular disease and cancer in later life: systematic review and meta-analysis. *British Medical Journal* **335** (7627), 974–978.
- Chai A. H., Wong T., Mak H.-L., et al. (2008) Prevalence and associated risk factors of retention of urine after caesarean section. *International Urogynaecology Journal* **19** (4), 537–542.
- Cherian D., Arianayagam M. & Rashid P. (2008) Symptomatic urinary stone disease in pregnancy. *Australian and New Zealand Journal of Obstetrics and Gynaecology* **48** (2), 1–27.
- de Leeuw J. W., de Wit C., Kuijken J. P. J. A. & Bruinse H. W. (2008) Mediolateral episiotomy reduces the risk for anal sphincter injury during operative vaginal delivery. *British Journal of Obstetrics and Gynaecology* **115** (1), 104–108.
- Dietz H. P. & Simpson J. M. (2007) Does delayed child-bearing increase the risk of levator injury in labour? *Australian and New Zealand Journal of Obstetrics and Gynaecology* **47** (6), 491–495.
- Duckett R. A. J., Grapsas P., Eaton M. & Basu M. (2008) The effect of spinal anaesthesia on urethral function. *International Urogynaecology Journal* **19** (2), 257–260.
- Ee C. C., Manheimer E., Pirotta M. V. & White A. R. (2008) Acupuncture for pelvic and back pain in pregnancy: a systematic review. *American Journal of Obstetrics and Gynecology* **198** (3), 254–259.
- Ekéus C., Nilsson E. & Gottvall K. (2008) Increasing incidence of anal sphincter tears among primiparas in Sweden: a population-based register study. *Acta Obstetricia et Gynecologica Scandanavica* **87** (5), 564–573.
- Elden H. H., Hagberg H., Fagevik Olsen M., Ladfors L. & Ostgaard H. C. (2008) Regression of pelvic girdle pain after delivery: follow-up of a randomised single blind controlled trial with different treatment modalities. *Acta Obstetricia et Gynecologica Scandanavica* **87** (2), 201–208.
- Gerdin E., Sverrisdottir G., Badi A., Carlsson B. & Graf W. (2007) The role of maternal age and episiotomy in the risk of anal sphincter tears during childbirth. *Australian and New Zealand Journal of Obstetrics and Gynaecology* **47** (4), 286–290.
- Gottvall K., Allebeck P. & Ekéus C. (2007) Risk factors for anal sphincter tears: the importance of maternal position at birth. *British Journal of Obstetrics and Gynaecology* **114** (10), 1266–1272.
- Hegaard H. K., Hedegaard M., Damm P., et al. (2008) Leisure time physical activity is associated with a reduced risk of preterm delivery. *American Journal of Obstetrics and Gynecology* **198** (2), 180–182.
- Hill K., Thomas K., AbouZahr C., et al. on behalf of the Maternal Mortality Working Group (2007) Estimates of maternal mortality worldwide between 1990 and 2005: an assessment of available data. *Lancet* **370** (9595), 1311–1319.
- Hoddinott P., Tappin D. & Wright C. (2008) Clinical review: breast feeding. *British Medical Journal* **336** (7649), 881–887.
- Ickovics J., Kershaw T. S., Westdahl C., et al. (2007) Group prenatal care and perinatal outcomes: a randomized controlled trial. *Obstetrical and Gynecological Survey* **62** (12), 766–767.
- Kalus S. M., Kornman L. H. & Quinlivan J. A. (2008) Managing back pain in pregnancy using a support garment: a randomised trial. *British Journal of Obstetrics and Gynaecology* **115** (1), 68–75.
- Kindberg S., Stehouwer M., Hvidman L. & Henriksen T. B. (2008) Postpartum perineal repair performed by midwives: a randomised trial comparing two suture techniques leaving the skin unsutured. *British Journal of Obstetrics and Gynaecology* **115** (4), 472–479.
- Madsen M., Jørgensen T., Jensen M. L., et al. (2007) Leisure time physical exercise during

- pregnancy and the risk of miscarriage: a study within the Danish National Birth Cohort. *British Journal of Obstetrics and Gynaecology* **114** (11), 1419–1426.
- Magnussen E. B., Vatten L. J., Lund-Nilsen T. I., et al. (2007) Pre-pregnancy cardiovascular risk factors as predictors of pre-eclampsia: population based cohort study. *British Medical Journal* **335** (7627), 978–982.
- Manheimer E., Zhang G., Udoff L., et al. (2008) Effects of acupuncture on rates of pregnancy and live birth among women undergoing in vitro fertilisation: systematic review and meta-analysis. *British Medical Journal* **336** (7643), 545–549.
- Molander P., Väyrynen T., Paavonen J., Lepistö A. & Luukkonen P. (2007) Outcome of primary repair of obstetric anal sphincter rupture using the overlap technique. *Acta Obstetricia et Gynecologica Scandanavica* **86** (12), 1458–1462.
- Prager M., Andersson K. L., Stephansson O., Marchionni M. & Marions L. (2008) The incidence of obstetric anal sphincter rupture in primiparous women: a comparison between two European delivery settings. *Acta Obstetricia et Gynecologica Scandanavica* **87** (2), 209–215.
- Richens Y. (2008) Tackling maternal obesity: suggestions for midwives. *British Journal of Midwifery* **16** (1), 14–19.
- Rodriguez A., Arenas E. A., Osorio A. L., Mendez O. & Zuleta J. J. (2008) Selective vs routine midline episiotomy for the prevention of third- or fourth-degree lacerations in nulliparous women. *American Journal of Obstetrics and Gynecology* **198** (3), 285–286.
- Serati M., Salvatore S., Khullar V., et al. (2008) Prospective study to assess risk factors for pelvic floor dysfunction after delivery. *Acta Obstetricia et Gynecologica Scandanavica* **87** (3), 313–318.
- Smith G. C. S. & Fretts R. C. (2007) Stillbirth. *Lancet* **370** (9600), 1715–1725.
- Symon A. (2008) Third degree tears: the three-stage negligence test. *British Journal of Midwifery* **16** (3), 192–193.
- Terry R., Niven C., Dowding D., et al. (2007) ‘Should I come in now?’: a study of women’s early labour experiences. *British Journal of Midwifery* **15** (10), 604–609.
- Todman D. (2007) A history of caesarean section: from ancient world to the modern era. *Australian and New Zealand Journal of Obstetrics and Gynaecology* **47** (5), 357–361.
- Tomson T. & Hiilesmaa V. (2007) Pregnancy plus: epilepsy in pregnancy. *British Medical Journal* **335** (7623), 769–773.
- Villar J., Carroli G., Zavaleta N., et al. for the World Health Organization 2005 Global Survey on Maternal and Perinatal Health Research Group (2007) Maternal and neonatal risks and benefits associated with caesarean section: multicentre prospective study. *British Medical Journal* **335** (7628), 1025–1030.
- Ward V. B. (2007) Pregnancy plus: eating disorders in pregnancy. *British Medical Journal* **336** (7635), 93–96.
- Wiener A. & Rogers C. (2008) Antenatal classes: women can’t think beyond labour. *British Journal of Midwifery* **16** (2), 121–124.
- ## Gynaecology oncology
- Collaborative Group on Endometrial Studies of Ovarian Cancer (2008) Ovarian cancer and oral contraceptives: collaborative reanalysis of data from 45 epidemiological studies including 23 257 women with ovarian cancer and 87 303 controls. *Lancet* **371** (9609), 304–314.
- Early Breast Cancer Trialists’ Collaborative Group (2008) Adjuvant chemotherapy in oestrogen-receptor-poor breast cancer: patient-level meta-analysis of randomised trials. *Lancet* **371** (9606), 29–40.
- International Collaboration of Epidemiological Studies of Cervical Cancer (2007) Cervical cancer and hormonal contraceptives: collaborative reanalysis of individual data for 16 573 women with cervical cancer and 35 509 women without cervical cancer from 24 epidemiological studies. *Lancet* **370** (9599), 1609–1621.
- James R. M., Cruikshank M. E. & Siddiqi N. on behalf of the Guideline Development Group (2008) Management of cervical cancer: summary of SIGN guidelines. *British Medical Journal* **336** (7634), 41–43.
- Reeves G., Pirie K., Beral V., et al. for the Million Women Study Collaboration (2007) Cancer incidence and mortality in relation to body mass index in the Million Women Study: cohort study. *British Medical Journal* **335** (7630), 1134–1139.
- Skjeldestad F. E. & Hagen B. (2008) Long-term consequences of gynaecological cancer treatment on urinary incontinence: a population-based cross-sectional study. *Acta Obstetricia et Gynecologica Scandanavica* **87** (3), 469–475.
- ## Bladder dysfunction
- Al-Buheissi S., Khasriya R., Maraj B. H. & Malone-Lee J. (2008) A simple validated scale to measure urgency. *Journal of Urology* **179** (3), 1000–1005.

- Ansari M. S., Srivastava A., Kapoor R., et al. (2008) Biofeedback therapy and home pelvic floor exercises for lower urinary tract dysfunction after posterior urethral valve ablation. *Journal of Urology* **179** (2), 708–701.
- Ansquer Y., Fernandez P., Aimot S., et al. (2007) MRI urethrovesical junction mobility is associated with global pelvic floor laxity in female stress incontinence. *Acta Obstetricia et Gynecologica Scandanavica* **86** (10), 1243–1250.
- Dumoulin C., Peng Q., Stodkilde-Jorgensen H., Shishido K. & Constantinou C. (2007) Changes in levator ani anatomical configuration following physiotherapy in women with stress urinary incontinence. *Journal of Urology* **178** (3), 970–977.
- Ebbesen M. H., Hannestad Y., Midthjell K. & Hunskaar S. (2007) Diabetes and urinary incontinence – prevalence data from Norway. *Acta Obstetricia et Gynecologica Scandanavica* **86** (10), 1256–1262.
- Eliasson K., Edner A. & Mattson E. (2008) Urinary incontinence in very young and mostly nulliparous women with a history of regular organized high-impact trampoline training: occurrence and risk factors. *International Urogynaecology Journal* **19** (5), 687–696.
- Gaskin C. J., Hawkins M. T. & O'Connell B. (2007) The construct validity of brief continence self-assessment awareness. *Australian and New Zealand Continence Journal* **13** (3), 61–65.
- Holroyd-Leduc J. M., Tannenbaum C., Thorpe K. E. & Straus S. E. (2008) What type of urinary incontinence does this woman have? *Journal of the American Medical Association* **299** (12), 1446–1456.
- Hullfish K., Fenner D., Sorser S., et al. (2007) Postpartum depression, urge incontinence and overactive bladder syndrome: is there an association? *International Urogynaecology Journal* **18** (10), 1121–1126.
- Kennedy C. M., Nygaard I. E., Bradley C. S. & Galask R. P. (2007) Bladder and bowel symptoms among women with vulvar disease: are they universal? *Journal of Reproductive Medicine* **52** (12), 1073–1079.
- Latthe P. M., Foon R. & Khan K. (2008) Nonsurgical treatment of stress urinary incontinence (SUI): grading of evidence in systematic reviews. *British Journal of Obstetrics and Gynaecology* **115** (4), 435–444.
- Miller J. M., Sampselle C., Ashton-Miller J., Hong G. S. & DeLancey J. O. (2008) Clarification and confirmation of the knack manoeuvre: the effect of volitional pelvic floor muscle contraction to preempt expected stress incontinence. *International Urogynaecology Journal* **19** (6), 773–782.
- Moore K. H. & Sarma S. (2007) The changing face of detrusor overactivity therapy in Australia. *Australian and New Zealand Continence Journal* **13** (3), 90–94.
- Morris A. R., Westbrook J. I. & Moore K. H. (2008) A longitudinal study over 5 to 10 years of clinical outcomes in women with idiopathic detrusor overactivity. *British Journal of Obstetrics and Gynaecology* **115** (2), 239–246.
- O'Reilly B. A., Fynes M., Achtari C., et al. (2008) A prospective randomized double-blind controlled trial evaluating the effect of trans-sacral magnetic stimulation in women with overactive bladder. *International Urogynaecology Journal* **19** (4), 497–502.
- Peters K., Girdler B., Carrico D., Ibrahim I. & Diokna A. (2008) Painful bladder syndrome/interstitial cystitis and vulvodynia: a clinical correlation. *International Urogynaecology Journal* **19** (5), 665–669.
- Salvatore S., Serati M., Ghezzi F., et al. (2007) Efficacy of tolterodine in women with detrusor overactivity and anterior vaginal wall prolapse: is it the same? *British Journal of Obstetrics and Gynaecology* **114** (11), 1436–1438.
- Sieger R. J., Weatherall M., Abernethy D. A. & Hay-Smith E. (2007) Bladder and bowel dysfunction in multiple sclerosis: a review of treatment effectiveness. *Australian and New Zealand Continence Journal* **13** (3), 81–89.
- Smith M. D., Russell A. & Hodges P. W. (2008) Is there a relationship between parity, pregnancy, back pain and incontinence? *International Urogynaecology Journal* **19** (3), 205–211.
- Sønksen J., Ohl D. A., Bonde B., Læssøe L. & McGuire E. J. (2007) Transcutaneous mechanical nerve stimulation using perineal vibration: a novel method for the treatment of female stress urinary incontinence *Journal of Urology* **178** (5), 2025–2028.
- Subak L., Van Den Eeden S., Thom D., Creasman J. M. & Brown J. S. (2007) Urinary incontinence in women: direct costs of routine care. *American Journal of Obstetrics and Gynecology* **197** (6), 596–598.
- Townsend M. K., Danforth K. N., Rosner B., et al. (2008) Physical activity and incident urinary incontinence in middle-aged women. *Journal of Urology* **179** (3), 1012–1017.
- Westergren H., Söderberg M., Johansson B., et al. (2007) Pelvic floor sex steroid hormone receptors, distribution and expression in pre- and postmenopausal stress urinary incontinent women. *Acta Obstetricia et Gynecologica Scandanavica* **86** (11), 1371–1384.

- Wijma J., Weiss Potters A. E., Tinga D. J. & Aarnoudse J. G. (2008) The diagnostic strength of the 24-h pad test for self-reported symptoms of urinary incontinence in pregnancy and after childbirth. *International Urogynaecology Journal* **19** (4), 525–530.
- Zabihi N., Mourtzinos A., Maher M. G., Raz S. & Rodriguez L. V. (2008) Short-term results of bilateral S2–S4 sacral neuromodulation for the treatment of refractory interstitial cystitis, painful bladder syndrome, and chronic pelvic pain. *International Urogynaecology Journal* **19** (4), 553–557.

Gynaecology surgery

- Altman D., Granath F., Cnattingius S. & Falconer C. (2007) Hysterectomy and risk of stress-urinary-incontinence surgery: nationwide cohort study. *Lancet* **370** (9597), 1494–1499.
- Diez-Itza I., Aizpitarte I. & Becerro A. (2007) Risk factors for the recurrence of pelvic organ prolapse after vaginal surgery: a review at 5 years after surgery. *International Urogynaecology Journal* **18** (11), 1317–1324.
- Gabriel B., Farthmann J., Brinstrup B., et al. (2007) Surgical repair of posterior compartment prolapse: preliminary results of a novel transvaginal procedure using a four-armed polypropylene mesh with infracoccygeal and pararectal suspension. *Acta Obstetricia et Gynecologica Scandanavica* **86** (10), 1236–1242.
- Gerten K. A., Richter H. E., Wheeler T. L., et al. (2008) Intraabdominal pressure changes associated with lifting: implications for postoperative activity restrictions. *American Journal of Obstetrics and Gynecology* **198** (3), 306–307.
- Glavind K., Morup L., Madsen H. & Glavind J. (2007) A prospective, randomized, controlled trial comparing 3 hour and 24 hour post-operative removal of bladder catheter and vaginal pack following vaginal prolapse surgery. *Obstetrical and Gynecological Survey* **62** (12), 780–781.
- Haensel A., Naumann G., Höhle P., Rommens P. M. & Koelbl H. (2007) Osteomyelitis following a transobturator sling (TVT-O). *British Journal of Obstetrics and Gynaecology* **114** (12), 1577–1579.
- Jelovsek J. E., Barber M. D., Karram M. M., Walters M. D. & Paraiso M. F. R. (2008) Randomised trial of laparoscopic Burch colposuspension versus tension-free vaginal tape: long-term follow up. *British Journal of Obstetrics and Gynaecology* **115** (2), 219–225.
- Kulseng-Hanssen S., Husby H. & Schiotz H. A. (2008) Follow up of TVT operations in 1113 women with mixed urinary incontinence at 7 and 38 months. *International Urogynaecology Journal* **19** (3), 391–396.
- Mallett V. T., Brubaker L., Stoddard A. M., et al. (2008) The expectations of patients who undergo surgery for stress incontinence. *American Journal of Obstetrics and Gynecology* **198** (3), 308–310.
- Poza J. L., Pla F., Sabadell J., et al. (2008) Trans-obturator suburethral tape for female stress incontinence: a cohort of 254 women with 1-year to 2-year follow-up. *Acta Obstetricia et Gynecologica Scandanavica* **87** (2), 232–239.
- Robert M., Soraisham A. & Sauve R. (2008) Postoperative urinary incontinence after total abdominal hysterectomy or supracervical hysterectomy: a metaanalysis. *American Journal of Obstetrics and Gynecology* **198** (3), 264–265.
- Ward K. L., Hilton P. & the UK and Ireland TVT Trial Group (2008) Tension-free vaginal tape versus colposuspension for primary urodynamic stress incontinence: 5-year follow up. *British Journal of Obstetrics and Gynaecology* **115** (2), 226–233.

Gynaecology

- Doshani A., Teo R. E. C., Mayne C. J. & Tincello D. G. (2007) Uterine prolapse. *British Medical Journal* **335** (7624), 819–823.
- Heiss G., Wallace R., Anderson G. L., et al. for the Women's Health Initiative Investigators (2008) Health risks and benefits 3 years after stopping randomized treatment with estrogen and progestin. *Journal of the American Medical Association* **299** (9), 1036–1045.
- Komesu Y. M., Rogers R. G., Rode M. A., et al. (2007) Pelvic floor symptom changes in pessary users. *American Journal of Obstetrics and Gynecology* **197** (6), 620–622.
- Nelson H. D. (2008) Menopause. *Lancet* **371** (9614), 760–770.
- Rogers R. G., Leeman L. M., Migliaccio L. & Albers L. L. (2008) Does the severity of spontaneous genital tract trauma affect postpartum pelvic floor function? *International Urogynaecology Journal* **19** (2), 429–435.
- Sewell C. A., Chang E. C. & Sultana C. J. (2007) Prevalence of genital prolapse in 3 ethnic groups. *Journal of Reproductive Medicine* **52** (9), 769–774.
- Spence D. & Melville C. (2007) Clinical review: vaginal discharge. *British Medical Journal* **335** (7630), 1147–1151.
- Srikrishna S., Robinson D., Cardozo L. & Gonzalez J. (2008) Is there a difference in

- patient quality of life evaluation in pelvic organ prolapse? *International Urogynaecology Journal* **19** (4), 517–520.
- Toozs-Hobson P., Balmforth J., Cardozo L., Khullar V. & Athanasiou S. (2008) The effect of mode of delivery on pelvic floor functional anatomy. *International Urogynaecology Journal* **19** (3), 407–416.
- Yonkers K. A., O'Brien P. M. S. & Eriksson E. (2008) Premenstrual syndrome. *Lancet* **371** (9619), 1200–1210.

Bowel dysfunction

- Altman D., Flaconer C., Roosner S. & Melin I. (2007) The risk of anal incontinence in obese women. *International Urogynaecology Journal* **18** (11), 1283–1289.
- Bradley C. S., Richter H. E., Gutman R. E., et al. (2007) Risk factors for sonographic internal anal sphincter gaps 6 to 12 months after delivery complicated by anal sphincter tear. *American Journal of Obstetrics and Gynecology* **197** (3), 310–311.
- Chiarelli P. (2008) The management of constipation using physical therapies including biofeedback. *Australian and New Zealand Continence Journal* **14** (1), 6–13.
- Handa V. L., Zyczynski H. M., Burgio K. L., et al. (2007) The impact of fecal and urinary incontinence on quality of life 6 months after childbirth. *American Journal of Obstetrics and Gynecology* **197** (6), 636–637.
- Morgan D. M., DeLancey J. O. L., Guire K. E. & Fenner D. E. (2007) Symptoms of anal incontinence and difficult defecation among women with prolapse and a matched control cohort. *American Journal of Obstetrics and Gynecology* **197** (5), 509–510.
- Mous M., Muller S. A. & de Leeuw J. W. (2008) Long-term effects of anal sphincter rupture during vaginal delivery: faecal incontinence and sexual complaints. *British Journal of Obstetrics and Gynaecology* **115** (2), 234–238.
- O'Boyle A. L., O'Boyle J. D., Magann E. F., et al. (2008) Anorectal symptoms in pregnancy and the postpartum period. *Journal of Reproductive Medicine* **53** (3), 151–155.
- Pretlove S. J., Thompson P. J., Toozs-Hobson P. M., Radley S. & Khan K. S. (2008) Does the mode of delivery predispose women to anal incontinence in the first year postpartum? A comparative systematic review. *British Journal of Obstetrics and Gynaecology* **115** (4), 421–434.
- Scheer I., Andrews V., Thakar R. & Sultan A. H. (2008) Urinary incontinence after obstetric anal sphincter injuries (OASIS) – is there a

relationship? *International Urogynaecology Journal* **19** (2), 179–183.

Sexual dysfunction

- Boij C., Matthiesen L., Krantz M. & Roland B. (2007) Sexual function and wellbeing after obstetric anal sphincter injury. *British Journal of Midwifery* **15** (11), 684–688.
- Pauls R. N., Silva W. A., Rooney C. M., et al. (2007) Sexual function after vaginal surgery for pelvic organ prolapse and urinary incontinence. *American Journal of Obstetrics and Gynecology* **197** (6), 622–625.
- Zahariou A. G., Karamouti M. V. & Papaioannou P. D. (2008) Pelvic floor muscle training improves sexual function of women with stress urinary incontinence. *International Urogynaecology Journal* **19** (3), 401–406.

Male health

- Niewoehner C. B. & Schorer A. E. (2008) Gynaecomastia and breast cancer in men. *British Medical Journal* **336** (7646), 709–713.

Miscellaneous

- Bolland M. J., Barber P. A., Doughty R. N., et al. (2008) Vascular events in healthy older women receiving calcium supplementation: randomised controlled trial. *British Medical Journal* **336** (7638), 262–266.
- Ellsberg M., Jansen H. A. F. M., Heise L., Watts C. H. & Garcia-Moreno C. on behalf of the WHO Multi-country Study on Women's Health and Domestic Violence against Women Study Team (2008) Intimate partner violence and women's physical and mental health in the WHO multi-country study on women's health and domestic violence: an observational study. *Lancet* **371** (9619), 1165–1172.
- Hilton P. (2008) Long-term follow-up studies in pelvic floor dysfunction: the Holy Grail or a realistic aim? *British Journal of Obstetrics and Gynaecology* **115** (2), 135–143.
- Mackey D. C., Lui L.-Y., Cawthorn P. M., et al. for the Study of Osteoporotic Fractures and Osteoporotic Fractures in Men Study Research Groups (2007) High-trauma fractures and low bone mineral density in older women and men. *Journal of the American Medical Association* **298** (20), 2381–2388.
- Plante A. F. & Kamm M. A. (2008) Life events in patients with vulvodynia. *British Journal of Obstetrics and Gynaecology* **115** (4), 509–514.
- Richens Y. (2007) Complications of catheterization. *British Journal of Midwifery* **15** (10), 622–625.

- Robbins J., Aragaki A. K., Kooperberg C., *et al.* (2007) Factors associated with 5-year risk of hip fracture in postmenopausal women. *Journal of the American Medical Association* **298** (20), 2389–2398.
- Tu F. F., Holt J., Gonzales J. & Fitzgerald C. M. (2008) Physical therapy evaluation of patients with chronic pelvic pain: a controlled study. *American Journal of Obstetrics and Gynecology* **198** (3), 272–275.